

*Pharmacochemistry*

## Study of Poly[Oxy-1-Carboxy-2-(3,4-Dihydroxyphenyl) Ethylene] from *Symphytum asperum*, *S. caucasicum*, *S. officinale*, *Anchusa italica* by Circular Dichroism

Vakhtang Barbakadze\*, Maia Merlani\*, Lela Amiranashvili\*,  
Lali Gogilashvili\*, Karen Mulkijanyan\*

\* I. Kutateladze Institute of Pharmacochemistry, Tbilisi State Medical University

(Presented by Academy Member Devi Ugrehelidze)

**ABSTRACT.** Comparative study of chirality of two carbon atoms C1 and C2 of poly[oxy-1-carboxy-2-(3,4-dihydroxyphenyl)ethylene] (POCDPE) preparations from different species of Boraginaceae family - *Symphytum asperum*, *S. caucasicum*, *S. officinale*, *Anchusa italica* revealed that they have one and the same absolute configuration in all preparations. It has been established that chiral atoms of POCDPE have either (1*R*,2*R*) or (1*S*,2*S*) configurations but not (1*R*,2*S*) and (1*S*,2*R*) configurations and consequently the denomination of the polymer is poly[oxy-(1*R*)-1-carboxy-(2*R*)-2-(3,4-dihydroxyphenyl)ethylene] or poly[oxy-(1*S*)-1-carboxy-(2*S*)-2-(3,4-dihydroxyphenyl)ethylene]. © 2012 Bull. Georg. Natl. Acad. Sci.

**Key words:** poly[3-(3,4-dihydroxyphenyl)glyceric acid], poly[oxy-1-carboxy-2-(3,4-dihydroxyphenyl)ethylene], 3-(3,4-dihydroxyphenyl)glyceric acid, *Symphytum asperum*, *S. caucasicum*, *S. officinale*, *Anchusa italica*, circular dichroism.

In previous papers we have reported about the isolation and identification of biologically active poly[3-(3,4-dihydroxyphenyl)glyceric acid] (PDPGA) that is poly[oxy-1-carboxy-2-(3,4-dihydroxyphenyl)ethylene] (POCDPE) from *Symphytum asperum* – POCDPE-SA, *S. caucasicum* – POCDPE-SC, *S. officinale* – POCDPE-SO and *Anchusa italica* – POCDPE-AI, respectively [1-8] (Fig. 1). The repeating unit of POCDPE – 3-(3,4-dihydroxyphenyl)glyceric acid residue contains two asymmetric carbon atoms C1 and C2 (Fig. 2), but we had not yet any information concerning the chirality of these centers.

Besides, the enantiomers of 3-(3,4-dihydroxyphenyl)glyceric acid – the monomer of PDPGA – (+)-(2*R*,3*S*)-2,3-dihydroxy-3-(3,4-dihydroxyphenyl)propionic acid [(2*R*,3*S*)-DDPPA] (Fig. 3) and (–)-(2*S*,3*R*)-2,3-dihydroxy-3-(3,4-dihydroxyphenyl)propionic acid [(2*S*,3*R*)-DDPPA] (Fig. 4) were synthesized from *trans*-caffeic acid [9].

**Results and Discussion.** Circular dichroism (CD) spectra of (2*R*,3*S*)-DDPPA and (2*S*,3*R*)-DDPPA are given in Figs. 5 and 6, respectively.

CD spectra show positive Cotton effect at the wavelength 195 nm and negative one at the wave-




Fig. 1. Poly[oxy-1-carboxy-2-(3,4-dihydroxyphenyl)ethylene] (POCDPE).




Fig. 2. Repeating unit of POCDPE-SA, POCDPE-SC, POCDPE-SO and POCDPE-AI.




Fig. 3. (+)-(2*R*,3*S*)-2,3-dihydroxy-3-(3,4-dihydroxyphenyl)-propionic acid [(2*R*,3*S*)-DDPPA].




Fig. 4. (-)-(2*S*,3*R*)-2,3-dihydroxy-3-(3,4-dihydroxyphenyl)-propionic acid [(2*S*,3*R*)-DDPPA].

lengths 209, 228 and 275 nm for (2*R*,3*S*)-DDPPA (Fig. 5) and vice versa for (2*S*,3*R*)-DDPPA (Fig. 6). Thus, Cotton effects in CD spectra of (2*R*,3*S*)-DDPPA and (2*S*,3*R*)-DDPPA have opposite profiles (Fig. 7).

The CD spectra of POCDPE-SA, POCDPE-SC, POCDPE-SO and POCDPE-AI have similar profiles and show positive Cotton effects at 194, 214, 280, 286

nm and a negative one at 204, 236 nm (Figs. 8 and 9). These data confirm that two chiral carbon atoms C1 and C2 of the repeating unit of POCDPE-SA, POCDPE-SC, POCDPE-SO and POCDPE-AI (Fig. 2) have one and the same absolute configuration.

Cotton effects in CD spectra of (2*R*,3*S*)-DDPPA and (2*S*,3*R*)-DDPPA are shifted about 5-11 nm




Fig. 5. CD spectrum of (2*R*,3*S*)-DDPPA.




Fig. 6. CD spectrum of (2*S*,3*R*)-DDPPA.




Fig. 7. Comparison of CD spectra of (2*R*,3*S*)-DDPPA (---) and (2*S*,3*R*)-DDPPA (—)

towards lower wavelengths vs Cotton effects in CD spectra of POCDPE-SA, POCDPE-SC, POCDPE-SO and POCDPE-AI.

Cotton effects at 195, 209, 228 and 275 nm in CD spectra of monomers – (2*R*,3*S*)-DDPPA and (2*S*,3*R*)-DDPPA correspond to Cotton effects at the 204, 214, 236 and 286 nm in CD spectra of POCDPE-SA, POCDPE-SC, POCDPE-SO and POCDPE-AI. Besides, as the profiles of CD spectra of polymers (Figs. 8 and 9) do not coincide with CD spectra of (2*R*,3*S*)-DDPPA and (2*S*,3*R*)-DDPPA (Fig. 7) we can exclude (1*R*,2*S*) and (1*S*,2*R*) configurations for chiral atoms C1 and C2 in the repeating units of POCDPE-SA, POCDPE-SC, POCDPE-SO and POCDPE-AI (Fig. 2). Hence, we can suppose that C1 and C2 atoms have either (1*R*,2*R*) or (1*S*,2*S*) configurations and consequently denomination of polymer is poly[oxy-(1*R*)-1-carboxy-(2*R*)-2-(3,4-dihydroxyphenyl)ethylene] or poly[oxy-(1*S*)-1-carboxy-(2*S*)-2-(3,4-dihydroxyphenyl)ethylene].

Establishment of the absolute configuration of these C atoms is a topic for further research.

**Materials and Methods.** CD spectra were performed on a Jasco J-715 instrument (Jasco Co, Tokyo,




Fig. 8. CD spectrum of POCDPE-SA.




Fig. 9. CD spectra of POCDPE-SC(1), POCDPE-SO(2) and POCDPE-AI(3).

Japan) equipped with Peltier temperature control system. For all measurements, 1 mm path length quartz cells, 1 nm bandwidth, 0.2 nm resolution, 1 s response and a scan speed of 50 nm/min for each spectrum were used.

CD spectrum of (2*R*,3*S*)-DDPPA ( $C = 0.56 \times 10^{-3}$  M,  $H_2O$ ):  $\Delta\epsilon_{275} = -1$ ,  $\Delta\epsilon_{228} = -5$ ,  $\Delta\epsilon_{209} = -10$ ,  $\Delta\epsilon_{195} = +5$ .

CD spectrum of (2*S*,3*R*)-DDPPA ( $C = 0.5 \times 10^{-3}$  M,  $H_2O$ ):  $\Delta\epsilon_{275} = -0.4$ ,  $\Delta\epsilon_{228} = +1.8$ ,  $\Delta\epsilon_{209} = +4.8$ ,  $\Delta\epsilon_{195} = -16$ .

CD spectrum of POCDPE-SA ( $C = 1.2$  mg/10 ml,  $H_2O$ ):  $\Delta\epsilon_{286} = +3$ ,  $\Delta\epsilon_{236} = -2$ ,  $\Delta\epsilon_{214} = +50$ ,  $\Delta\epsilon_{204} = -28$ ,  $\Delta\epsilon_{194} = +4$ .

## ფარმაკოქიმია

## *Symphytum asperum*-ის, *S. caucasicum*-ის, *S. officinale*-ის, *Anchusa italica*-ს პოლი[ოქსი-1-კარბოქსი-2-(3,4-დიჰიდროქსიფენილ)ეთილენის] შესწავლა წრიული დიქროიზმის მეთოდით

ვ. ბარბაქაძე\*, მ. მერლანი\*, ლ. ამირანაშვილი\*, ლ. გოგილაშვილი\*,  
კ. მულკიჯანიანი\*

\* *Tbilisis saxel mwi fo samedicino universiteti, i.qutaTel azis farmakoqimii instituti, Tbilisi*

(წარმოდგენილია აკადემიის წევრის დ. უგრეხელიძის მიერ)

Boraginaceae-ს ოჯახის სხვადასხვა სახეობების (*Symphytum asperum*, *S. caucasicum*, *S. officinale* და *Anchusa italica*) პოლი[ოქსი-1-კარბოქსი-2-(3,4-დიჰიდროქსიფენილ)ეთილენის] (პოკდფე) პრეპარატების ორი ნაზშირბად ატომის C1 და C2 ქირალობის შედარებითმა შესწავლამ აჩვენა, რომ ყველა პრეპარატში მათ აქვთ ერთი და იგივე აბსოლუტური კონფიგურაცია. დადგენილია, რომ პოკდფე-ს ქირალურ ატომებს აქვთ ან (1R,2R) ან (1S,2S) კონფიგურაციები, მაგრამ არა (1R,2S) და (1S,2R) კონფიგურაციები და აქედან გამომდინარე პოლიმერის სახელწოდება შეიძლება იყოს პოლი[ოქსი-(1R)-1-კარბოქსი-(2R)-2-(3,4-დიჰიდროქსიფენილ)ეთილენი] ან პოლი[ოქსი-(1S)-1-კარბოქსი-(2S)-2-(3,4-დიჰიდროქსიფენილ)ეთილენი].

## REFERENCES

1. V.V.Barbakadze, E.P.Kemertelidze, A.S.Shashkov, A.I.Usov (2000), *Mendeleev Commun.*, **10**, 4: 148-149.
2. C.M.Barthomeuf, E.Debiton, V.V.Barbakadze, E.P.Kemertelidze (2001), *J. Agric. Food Chem.*, **49**, 8: 3942-3946.
3. V.V.Barbakadze, E.P.Kemertelidze, I.L.Targamadze, et al. (2002), *Russ. J. Bioorg. Chem.*, **28**, 4: 326-330.
4. V.Barbakadze, E.Kemertelidze, I.Targamadze, et al. (2005), *Molecules*, **10**, 9: 1135-1144.
5. V.V.Barbakadze, E.P.Kemertelidze, K.G.Mulkijanyan, et al. (2007), *Pharm. Chem. J.*, **41**, 1: 14-16.
6. V.Barbakadze, A.J.J. van den Berg, C.J.Beukelman, et al. (2009), *Chem. Nat. Compds.*, **45**, 1: 6-10.
7. V. Barbakadze, L.Gogilashvili, L.Amiranashvili, et al. (2010), *Nat. Prod. Commun.*, **5**, 7: 1091-1095.
8. V.Barbakadze, K.Mulkijanyan, M.Merlani, et al. (2008), *Bull. Georg. Natl. Acad. Sci.*, **2**, 3, 108-112.
9. M.Merlani, V.Barbakadze, L.Amiranashvili, et al. (2010), *Chirality*, **22**, 8: 717-725.

Received December, 2011